



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE CHORD COMMON TO A PARABOLA AND THE CIRCLE OF CURVATURE AT ANY POINT.

By PROF. R. H. GRAVES, Chapel Hill, N. C.

It is known that if a circle meet a parabola in four points, the sum of the distances of the points on one side of the axis from it is equal to the sum of the distances of the points on the other side from it. If three of the points are coincident, the circle becomes the circle of curvature, and the distance of the three coincident points (P) from the axis is one-third of that of the fourth point from the axis.

Hence the common chord of the circle and parabola is divided by the axis in the ratio 1 : 3. But the shorter segment of the chord is equal to the tangent at P , since they are equally inclined to the axis. Therefore the chord is equal to four times the tangent. Let $y^2 = 4ax$ be the equation to the parabola, and (x', y') the co-ordinates of P . Then

$$y - y' = -\frac{2a}{y'}(x - x'), \quad \text{or} \quad yy' + 2ax - \frac{3}{2}y'^2 = 0,$$

is the equation to the chord.

Differentiating with respect to y' , $y = 3y'$; hence $y^2 = -12ax$ is the envelope of the chord. Also, from relation $y = 3y'$, it follows that the longer segment of the chord is equal to the corresponding tangent of the parabola, $y^2 = -12ax$.

The point P , and the point where the chord prolonged touches $y^2 = -12ax$, are harmonic conjugates with respect to the points where it meets the axis and the tangent at the common vertex of the parabolas.

The tangent at the end of the *latus rectum* of $y^2 = -12ax$ is normal to $y^2 = 4ax$ at the end of its *latus rectum*, and therefore touches its evolute. The chord is then a diameter of the circle of curvature, and is bisected by its point of contact with the evolute.

Hence the radius of curvature = twice the normal = $4a\sqrt{2}$, which agrees with a known result.